



basic education

Department:
Basic Education
REPUBLIC OF SOUTH AFRICA

**NATIONAL
SENIOR CERTIFICATE**

GRADE 12

TECHNICAL MATHEMATICS P2

MARKING GUIDELINES

EXEMPLAR 2018

MARKS: 150


These marking guidelines consist of 14 pages

QUESTION 1

| | | |
|------------|--|--|
| <p>1.1</p> | $AD = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$ $= \sqrt{(5-1)^2 + (-2-4)^2}$ $= \sqrt{52}$ $= 2\sqrt{13}$ | <p>✓ Substitution in correct formula ✓ Simplification ✓ $2\sqrt{13}$ simplified surd (3)</p> |
| <p>1.2</p> | $M\left(\frac{x_1 + x_2}{2}; \frac{y_1 + y_2}{2}\right)$ $M\left(\frac{1+5}{2}; \frac{4-2}{2}\right)$ $M(3; 1)$ | <p>✓ Substitution ✓ M(3; 1) (2)</p> |
| <p>1.3</p> | $m_{AB} = \frac{y_2 - y_1}{x_2 - x_1}$ $= \frac{1-4}{-3-1}$ $= \frac{3}{4}$ <p>AB MC</p> $m_{MC} = \frac{3}{4}$ $y - y_1 = m(x - x_1) \quad \text{or} \quad y = mx + c$ $y - 1 = \frac{3}{4}(x - 3) \quad \quad \quad 1 = \frac{3}{4}(3) + c$ $-3x + 4y + 5 = 0 \quad \quad \quad c = \frac{4-9}{4} = -\frac{5}{4}$ $\quad \quad \quad \text{or } 3x + -4y - 5 = 0 \quad \quad \quad y = \frac{3}{4}x - \frac{5}{4}$ $\quad \quad \quad \quad \quad \quad \quad \quad \quad -3x + 4y + 5 = 0$ | <p>✓ Substitution in correct formula ✓ Gradient of MC ✓ Substitution in correct formula ✓ Simplification ✓ Correct answer in correct form (5)</p> |
| <p>1.4</p> | $\tan \alpha = \frac{3}{4}$ $\alpha \approx 36,87^\circ$ | <p>✓ $\tan \alpha = \frac{3}{4}$ ✓ $36,87^\circ$ (2)</p> |

| | | |
|------------|--|--|
| <p>1.5</p> | $m_{AD} = \frac{y_2 - y_1}{x_2 - x_1}$ $= \frac{-2 - 4}{5 - 1}$ $= \frac{-3}{2}$ $\tan \beta = \frac{-3}{2}$ $\beta \approx 123,69^\circ$ $\hat{B}AD = 123,69^\circ - 36,87^\circ \quad \text{ext } \angle \text{ of } \Delta$ $= 86,82^\circ$ | $\checkmark \frac{-3}{2}$ $\checkmark \beta = 123,69^\circ$ $\checkmark 123,69^\circ - 36,87^\circ$ $\checkmark 86,82^\circ$ <p style="text-align: right;">(4)</p> |
| | | <p>[16]</p> |

QUESTION 2

| | | |
|--------------|---|---|
| <p>2.1.1</p> | $x^2 + y^2 = r^2$ $r^2 = (10)^2 + (-4)^2$ $\therefore r = \sqrt{116} = 2\sqrt{29}$ | <p>✓ Substitution.</p> <p>✓ $\sqrt{116} = 2\sqrt{29}$</p> <p>(2)</p> |
| <p>2.1.2</p> | $m_{OH} = m_{OF} = \frac{0 - (-4)}{0 - 10}$ $= -\frac{4}{10}$ $= -\frac{2}{5}$ | <p>✓ Substitution.</p> <p>✓ $-\frac{2}{5}$</p> <p>(2)</p> |
| <p>2.1.3</p> | $m_{GH} = \frac{5}{2} \quad (m_{GH} \times m_{OH} = -1)$ <p>By symmetry H(-10; 4)</p> $y - y_1 = m(x - x_1) \quad \text{or} \quad y = mx + c$ $y - 4 = \frac{5}{2}(x + 10) \quad \quad \quad 4 = \frac{5}{2}(-10) + c$ $y = \frac{5}{2}x + 29 \quad \quad \quad c = 29$ $y = \frac{5}{2}x + 29$ | <p>✓ m_{GH}</p> <p>✓ H(-10; 4)</p> <p>✓ Substitution of H</p> <p>✓ Equation</p> <p>(4)</p> |
| <p>2.2</p> |  | <p>✓ Both x-intercepts</p> <p>✓ Both y-intercepts</p> <p>✓ Shape</p> <p>(3)</p> |
| | | <p>[11]</p> |

QUESTION 3

| | | |
|-------|---|---|
| 3.1.1 | $OP^2 = (12)^2 + (-5)^2$ $\therefore OP = 13 \text{ units}$ | $\checkmark OP = 13 \text{ units}$ <p style="text-align: right;">(1)</p> |
| 3.1.2 | $5 \cot \theta - 13 \cos \theta$ $= 5 \left(\frac{12}{-5} \right) - 13 \left(\frac{12}{13} \right)$ $= -24$ | $\checkmark \frac{12}{-5} \quad \checkmark \frac{12}{13}$ $\checkmark \text{Simplification}$ <p style="text-align: right;">(3)</p> |
| 3.1.3 | $\operatorname{cosec}^2 x - 1$ $= \left(\frac{13}{-5} \right)^2 - 1$ $= \frac{144}{25}$ <p style="text-align: center;"><i>or</i></p> $\operatorname{cosec}^2 x - 1 = \cot^2 x$ $= \left(\frac{12}{-5} \right)^2$ $= \frac{144}{25}$ | $\checkmark \left(\frac{13}{-5} \right)^2 \text{ or } \left(\frac{12}{-5} \right)^2$ $\checkmark \text{Simplification}$ <p style="text-align: right;">(2)</p> |
| 3.2 | $\sec(a - b)$ $= \sec(2,659 - 1,112)$ $= \sec 1,547$ $= \frac{1}{\cos 1,547}$ $\approx 42,03$ | $\checkmark \text{Substitution}$ $\checkmark \text{Reciprocal}$ $\checkmark \text{Simplification using radians}$ <p style="text-align: right;">(3)</p> |
| | | [9] |

QUESTION 4

| | | |
|------------|---|--|
| <p>4.1</p> | $\frac{\sin(360^\circ - x) \cdot \cos(180^\circ - x) \cdot \tan 120^\circ}{\cos^2 x \cdot \sin \frac{5\pi}{6}}$ $= \frac{(-\sin x) \cdot (-\cos x) \cdot (-\tan 60^\circ)}{\cos^2 x \cdot \sin \frac{\pi}{6}}$ $= \frac{-\sin x \cdot \cos x \cdot -\sqrt{3}}{\cos^2 x \cdot \frac{1}{2}}$ $= \frac{2\sqrt{3} \sin x}{\cos x}$ $= 2\sqrt{3} \tan x$ | <p>✓ $-\sin x$ ✓ $-\cos x$ ✓ $-\tan 60^\circ$</p> <p>✓ $\sin \frac{\pi}{6}$</p> <p>✓ $-\sqrt{3}$ and $\frac{1}{2}$</p> <p>✓ simplification</p> <p>✓ $\tan x$</p> <p>(7)</p> |
| <p>4.2</p> | <p>$\cos^2 3x$</p> | <p>✓ $\cos^2 3x$</p> <p>(1)</p> |
| <p>4.3</p> | <p>$LHS = \frac{\sin x}{\cos x} (\sin x)$</p> $= \frac{\sin^2 x}{\cos x}$ <p>$RHS = \sec x - \cos x$</p> $= \frac{1}{\cos x} - \cos x$ $= \frac{1 - \cos^2 x}{\cos x}$ $= \frac{\sin^2 x}{\cos x}$ <p>$= LHS$</p> | <p>✓ $\frac{\sin x}{\cos x}$</p> <p>✓ $\frac{1}{\cos x}$</p> <p>✓ $\frac{1 - \cos^2 x}{\cos x}$</p> <p>✓ $\sin^2 x$</p> <p>(4)</p> |
| <p>4.4</p> | <p>$\operatorname{cosec} 2x = 2,114$</p> $\sin 2x = \frac{1}{2,114}$ $2x = \sin^{-1}\left(\frac{1}{2,114}\right)$ $2x = \sin^{-1}\left(\frac{1}{2,114}\right)$ <p>$2x \approx 28,23^\circ$ or $2x \approx 180^\circ - 28,23^\circ$</p> <p>$x \approx 14,12^\circ$ or $x \approx 75,89^\circ$</p> | <p>✓ using reciprocal</p> <p>✓ inverse</p> <p>✓ value of x ✓ another value of x</p> <p>(4)</p> |
| | | <p>[16]</p> |

QUESTION 5

| | | |
|------------|---|--|
| <p>5.1</p> | $\sin 55^\circ = \frac{50}{AC}$ $AC = \frac{50}{\sin 55^\circ}$ $\approx 61 \text{ m}$ | <p>✓ Definition</p> <p>✓ AC subject</p> <p>✓ 61 m</p> <p style="text-align: right;">(3)</p> |
| <p>5.2</p> | <p>$AD \approx 61 \text{ m}$</p> $DC^2 = AC^2 + AD^2 - 2AC \cdot AD \cos 65^\circ$ $DC^2 = (61)^2 + (61)^2 - 2(61)(61) \cos 65^\circ$ <p>$DC \approx 66 \text{ m}$</p> | <p>✓ $AD \approx 61 \text{ m}$</p> <p>✓ Using cosine rule</p> <p>✓ Substitution</p> <p>✓ 66 m</p> <p style="text-align: right;">(4)</p> |
| <p>5.3</p> | $BD = \sqrt{AD^2 - AB^2}$ $= \sqrt{(61)^2 - (50)^2} \quad \text{OR} \quad \tan 55^\circ = \frac{50}{BD}$ $\approx 35 \text{ m} \quad \quad \quad = 35 \text{ m}$ $\therefore \text{area of } \triangle BDC = \frac{1}{2}(35)(66) \sin \hat{BDC} = 563$ $\sin \hat{BDC} = \frac{563}{\frac{1}{2}(35)(66)}$ $\therefore \hat{BDC} = \sin^{-1}\left(\frac{563}{\frac{1}{2}(35)(66)}\right)$ $= 29,17^\circ$ | <p>✓ Using Pythagoras theorem or tan</p> <p>✓ $BD = 35 \text{ m}$</p> <p>✓ Substitution in area formula</p> <p>✓ Area = 563</p> $\checkmark \sin \hat{BDC} = \frac{563}{\frac{1}{2}(35)(66)}$ <p>✓ Simplification</p> <p style="text-align: right;">(6)</p> |
| | | <p>[13]</p> |

QUESTION 6

| | | |
|------------|--|---|
| <p>6.1</p> | | <p>$f(x)$ ✓ x-intercepts ✓ y-intercept ✓ Shape</p> <p>$g(x)$ ✓ y-intercept ✓ Turning points ✓ Shape</p> <p>(6)</p> |
| <p>6.2</p> | <p>2</p> | <p>✓2 (1)</p> |
| <p>6.3</p> | <p>360°</p> | <p>✓ 360° (1)</p> |
| <p>6.4</p> | <p>$60^\circ < x < 240^\circ$</p> | <p>✓✓ Answer with correct notation (2)</p> |
| | | <p>[10]</p> |

QUESTION 7

| | | |
|--------|---|--|
| 7.1 | the angle in the alternate segment | ✓ Correct statement (1) |
| 7.2.1 | $\hat{R}_2 = \hat{S}_1 = 38^\circ$ tan – chord theorem | ✓ Statement ✓ Reason (2) |
| 7.2.2 | $\hat{M}_1 = 2\hat{R}_2 = 76^\circ$ \angle at center = 2 \angle at circum | ✓ Statement ✓ Reason (2) |
| 7.2.3. | $\hat{S}_2 = 90^\circ - 38^\circ = 52^\circ$ tan \perp radius | ✓ Statement ✓ Reason (2) |
| 7.2.4 | $\hat{R}_1 = 90^\circ$ \angle in a semi – circle $\hat{Q}_1 = \hat{S}_1 = 38^\circ$ tan– chord theorem $\therefore \hat{Q}_2 = 180^\circ - (17^\circ + 90^\circ + 38^\circ)$ sum of \angle 's of a Δ $= 35^\circ$ | ✓ Statement ✓ Reason ✓ Reason ✓ Statement ✓ Statement (5) |
| 7.2.5 | $\hat{R}_2 = 38^\circ$ and $\hat{P}_1 = 17^\circ$ $\therefore \hat{R}_2 \neq \hat{P}_1$ (alternate angles are not equal) | ✓ Alternate angles are not equal <i>or</i> $\hat{R}_2 \neq \hat{P}_1$ <i>or</i> $\hat{R}_2 = 38^\circ$ and $\hat{P}_1 = 17^\circ$ (1) |
| | | [13] |

QUESTION 8

| | | |
|-------|--|--|
| 8.1 | divides the other two sides proportionally | ✓ Answer (1) |
| 8.2.1 | $\frac{x}{8} = \frac{4}{10}$ $10x = 32$ $x = 3,2$ | ✓ Prop. ✓ Simplification ✓ Value of x (3) |
| 8.2.2 | RTSP is a parallelogram (both pairs of opposite sides of a quadrilateral are parallel) | ✓ Statement ✓ Reason (2) |
| 8.2.3 | $\frac{y}{9} = \frac{3,2}{8}$ Prop. Theorem ; TS MP $y = 3,6$ | ✓ Statement ✓ Reason ✓ Value of y (3) |
| 8.2.4 | $\frac{MR}{TS} = \frac{10}{4} = 2,5$ $\frac{RT}{SN} = \frac{9}{3,6} = 2,5$ $\frac{MT}{TN} = \frac{8}{3,2} = 2,5$ $\Delta MRT \parallel \Delta TSN$ sides of triangle are in the same proportion | ✓ Ratio ✓ Ratio ✓ Ratio ✓ Reason (4) |
| | | [13] |

QUESTION 9

| | | |
|-------------|--|---|
| 9.1 | $\hat{K}_1 = \hat{LHF}$ ext \angle of cyclic quad $= \hat{GFK}$ corres. angles LK // GF | ✓ Statement ✓ Reason ✓ Statement/Reason (3) |
| 9.2 | $\hat{MFG} = \hat{K}_1 = 104^\circ$ <i>corres\angles; FG \square KL</i> $\hat{G} + 104^\circ + 20^\circ = 180^\circ$ <i>angles of Δ</i> $\hat{G} = 56^\circ$ $\hat{MFG} = \hat{K}_1 = 104^\circ$ <i>corres. \angles; FG // KL</i> $\hat{G} + 104^\circ + 20^\circ = 180^\circ$ <i>angles of Δ</i> $\hat{G} = 56^\circ$ OR $\hat{MLK} + 104^\circ + 20^\circ = 180^\circ$ <i>angles of Δ \angles; FG // KL</i> $\hat{MLK} = 56^\circ$ $\hat{MLK} = \hat{G}$ <i>corresp \angles; FG // KL</i> $\hat{G} = 56^\circ$ | ✓ Statement ✓ Statement ✓ Reason ✓ Statement ✓ Statement ✓ Reason (3) |
| 9.3.1 | $\frac{10}{30} = \frac{12}{MG}$ (Prop. Theorem; KL // FG) $MG = \frac{360}{10} = 36 \text{ units}$ | ✓ Statement ✓ Reason ✓ 36 units (3) |
| 9.3.2 | $\hat{MHF} = \hat{GFM} = 104^\circ$ proved in question 9.1. \hat{M} is common $\hat{G} = \hat{F}_2$ sum of angles of a Δ $\Delta MFH \parallel \Delta MGF$ $\angle; \angle; \angle$ | ✓ Statement ✓ Statement and reason ✓ Reason (3) |
| 9.3.3 | $\Delta MFH \parallel \Delta MGF \parallel \Delta MLK$ | ✓ ΔMLK (1) |
| [13] | | |

QUESTION 10

| | | |
|--------|--|---|
| 10.1 | $x^2 - 4dh + 4h^2 = 0$ $x^2 - (4 \times 220 \text{ mm} \times 60 \text{ mm}) + 4(60 \text{ mm})^2 = 0$ $x^2 = \sqrt{67200 \text{ mm}^2}$ $\therefore x = 259,23 \text{ mm}$ | ✓Formula ✓Substitution. ✓Simplifying ✓Length (4) |
| 10.2.1 | $v = \pi Dn$ $= \pi(18 \text{ m})\left(\frac{225}{60 \text{ s}}\right)$ $\approx 212,06 \text{ m/s}$ | ✓Correct formula ✓Correct diameter ✓Substitution ✓212,06 m/s (4) |
| 10.2.2 | $\omega = 2\pi n$ $= 2\pi\left(\frac{225}{60 \text{ s}}\right)$ $\approx 23,56 \text{ rad/s}$ | ✓Correct formula ✓Substitution ✓23,56 rad/s (3) |
| 10.3.1 | $\hat{LBA} = 180^\circ - 70^\circ = 110^\circ$ co-int.angles; AK // BL | ✓Statement ✓Reason (2) |
| 10.3.2 | $\hat{KAD} = 360^\circ - 140^\circ = 220^\circ$ $\approx 3,84 \text{ rad}$ $s_1 = r\theta = 50(3,84)$ $\approx 192 \text{ cm}$ $s_2 = 48,8 \text{ cm}$ $\text{Length of belt} = 110 + 192 + 110 + 48,80$ $= 460,80 \text{ cm}$ | ✓Size of \hat{KAD} ✓Conversion to radians ✓Correct substitution in arc length formula ✓Arc length s_1 ✓Length of DE = 110 ✓Length of belt (6) |
| | | [19] |

QUESTION 11

| | | |
|---------------|---|---|
| <p>11.1.1</p> | <p>Volume of pyramid $= \frac{1}{3}(\text{area of base})(\text{height})$ $= \frac{1}{3}(4m \times 4m)(1,1m)$ $= 5,87m^3$ Volume of Cube $= (l)(b)(h)$ $= (4m)(4m)(4m)$ $= 64m^3$ Total Volume $= 5,87 m^3 + 64 m^3$ $\approx 69,87m^3$</p> | <p>✓ Substitution in correct formula ✓ Height ✓ $5,87m^3$ ✓ Substitution in correct formula ✓ $64m^3$ ✓ $69,87m^3$ (6)</p> |
| <p>11.1.2</p> | <p>total surface area $=$ surface area of cube base + surface area of pyramid $= 4(\text{side} \times \text{side}) + 4\left(\frac{1}{2} \times \text{base} \times \text{slant height}\right)$ $= 4(4m \times 4m) + 4\left(\frac{1}{2} \times 4m \times \sqrt{1,1+2^2} m\right)$ $= 64m^2 + 4(2m \times \sqrt{5,1} m)$ $= 82,07m^2$</p> | <p>✓ Slant height ✓ Correct substitution into area of a cube ✓ Correct substitution into area of pyramid ✓ Simplification ✓ Total area (5)</p> |
| <p>11.1.3</p> | <p>Cost of paint $= 82,07 \times R30,50$ $= R2503,14$</p> | <p>✓ $82,07 \times R30,50$ ✓ $R2503,14$ (2)</p> |

| | | |
|-------------|---|---|
| <p>11.2</p> | $A_T = a \left(\frac{o_1 + o_n}{2} + o_2 + o_2 + o_3 + \dots + o_{n-1} \right)$ $= 4 \left(\frac{6,2 + 2}{2} + y + 5,1 + 4,9 \right)$ $= 4(14,1 + y)$ $= 56,4 + 4y$ $\therefore 72 = 56,4 + 4y$ $\Rightarrow y = \frac{15,6}{4}$ $= 3,9m$ <p>OR</p> $A_T = a(m_1 + m_2 + m_3 + \dots + m_n)$ $72 = 4 \left(\frac{6,2 + y}{2} + \frac{y + 5,1}{2} + \frac{5,1 + 4,9}{2} + \frac{4,9 + 2}{2} \right)$ $18 = \frac{11,3 + 2y}{2} + 8,45$ $9,55 = \frac{11,3 + 2y}{2}$ $19,1 = 11,3 + 2y$ $\therefore y = 3,9m$ | <p>✓Formula</p> <p>✓Substitution</p> <p>✓Simplification</p> <p>✓ 3,9m</p> <p>OR</p> <p>✓Formula</p> <p>✓Substitution</p> <p>✓Simplification</p> <p>✓ 3,9m</p> <p>(4)</p> <p>[17]</p> |
|-------------|---|---|

TOTAL: 150