



basic education

Department:
Basic Education
REPUBLIC OF SOUTH AFRICA

**NATIONAL
SENIOR CERTIFICATE**

GRADE 12

AGRICULTURAL SCIENCES P1

NOVEMBER 2013

MEMORANDUM

MARKS: 150

This memorandum consists of 10 pages.

SECTION A**QUESTION 1**

1.1	1.1.1	D ✓✓	(10 x 2)	(20)
	1.1.2	B ✓✓		
	1.1.3	D ✓✓		
	1.1.4	C ✓✓		
	1.1.5	C ✓✓		
	1.1.6	A ✓✓		
	1.1.7	A ✓✓		
	1.1.8	B ✓✓		
	1.1.9	C ✓✓		
	1.1.10	B ✓✓		
1.2	1.2.1	B only ✓✓	(5 x 2)	(10)
	1.2.2	B only ✓✓		
	1.2.3	A only ✓✓		
	1.2.4	None ✓✓		
	1.2.5	A only ✓✓		
1.3	1.3.1	Small intestine/jejunum/ileum/duodenum ✓✓	(5 x 2)	(10)
	1.3.2	Carbohydrates/fats/lipids/oils ✓✓		
	1.3.3	Lobola ✓✓		
	1.3.4	Ovulation ✓✓		
	1.3.5	Quarantine ✓✓		
1.4	1.4.1	Amino acids/peptides ✓	(5 x 1)	(5)
	1.4.2	Endothermic/homoeothermic/warm blooded ✓		
	1.4.3	Indigenous ✓		
	1.4.4	Feedlot ✓		
	1.4.5	Blowflies ✓		

TOTAL SECTION A: 45

SECTION B**QUESTION 2: ANIMAL NUTRITION****2.1 Digestive system of a farm animal****2.1.1 Labelled parts**

- (a) D ✓ (1)
- (b) G ✓ (1)
- (c) C ✓ (1)

2.1.2 THREE adaptations of part C for absorption.

- It is a long thin tube that allows more nutrients to have contact with the walls and makes absorption easier ✓
- It has numerous folds hence a large surface area and makes more absorption possible ✓
- It has finger-like projections that enlarge the surface area for absorption ✓
- Villi with blood vessels/micro villi allow for easier absorption of nutrients into the bloodstream ✓
- Slow movement of food/muscle movement allows for more contact time with nutrients that will be absorbed ✓
- The presence of a single layer of columnar epithelial cells makes it easier for rapid absorption of nutrients ✓ (Any 3) (3)

2.1.3 Absorption of fat molecules

- The fat molecules are broken up into fatty acids and glycerol/ enzyme lipase breaks up fat molecules into simpler and soluble substances ✓
- Short chain fatty acids/ simpler soluble substances are directly absorbed ✓
- Passive absorption into the blood capillaries through diffusion ✓
- Fatty acids and glycerol are absorbed into the lacteal / lymph vessel ✓
- Long chain fatty acids are actively absorbed through carrier molecules ✓ (Any 2) (2)

2.2 Digestibility of feed**2.2.1 Digestibility co-efficient**

$$\begin{aligned} \text{DC} &= \frac{\text{DM intake (kg)} - \text{DM of manure (kg)} \times 100}{\text{DM intake(kg)}} \checkmark \\ &= \frac{8.8 \text{ kg} - 2.7 \text{ kg} \times 100}{8.8 \text{ kg}} \checkmark \end{aligned}$$

Or

$$= \frac{6.1 \times 100}{8.8 \text{ kg}} \checkmark$$

Or


One mark for evidence of substitution of values into the formula ✓
One mark for evidence of simplification of values ✓

And

$$= 69,32 \text{ Or } 69 \checkmark \% \checkmark \quad (5)$$

2.2.2 Implication of the value obtained in QUESTION 2.2.1

- 69,32 or 69% of the feed ✓
- was digested and absorbed ✓ **Or**
- 30,68 or 31% of the feed ✓
- was not digested and absorbed but excreted ✓ (Any 2) (2)

2.3 Balancing rations2.3.1 **Feed A (Maize):** 9 **20 ✓****Feed B: (OCM)**
(Oil cake meal) 38 **9 ✓**

Mix 20 parts of Feed A (Maize) with 9 parts of Feed B (OCM) **Or** 20:9 ✓ (4)

2.3.2 Percentage of maize

$$20 + 9 = 29 \checkmark$$

$$= \frac{20 \times 100}{29} \checkmark$$

$$= 68,97 \text{ Or } 69\% \checkmark \quad (3)$$

2.4 Nutrient deficiencies

- 2.4.1 Phosphorus/P ✓ (1)
- 2.4.2 Iron/Fe ✓ (1)
- 2.4.3 Iodine/I ✓ (1)

2.5 Comparing feeds**2.5.1 Nutritive ratio
Feed A:**

$$\text{TDN} = 8 + 50 + 22 = 80\% \checkmark \text{ Or } \text{DNNE} = 50 + 22 = 72\% \checkmark$$

$$\text{NR} = 1: \frac{\text{TDN}-\text{DP}}{\text{DP}} \checkmark \text{ Or } 1: \frac{\text{DNNE}'\text{s}}{\text{DP}} \checkmark$$

$$\text{NR} = 1: \frac{80-8}{8} \checkmark \text{ Or } 1: \frac{72}{8} \checkmark$$

$$\text{NR} = 1: 9 \checkmark \quad (4)$$

2.5.2 Feed for fattening animals with reason

- Feed A ✓ (1)
- Reason**
- NR is wide/(1:9) ✓
- Higher ratio of carbohydrates to protein ✓
- As carbohydrates are necessary for fattening ✓ (Any 2) (2)

2.5.3 Feed suited for young growing animals

- Feed B ✓ (1)
- Reason**
- NR is narrow/(1:5) ✓
- Higher ratio protein to carbohydrates ✓
- As protein is necessary for growth ✓ (Any 2) (2)

[35]**QUESTION 3: ANIMAL PRODUCTION****3.1 Animal behaviour**

- 3.1.1 E ✓ (1)
- 3.1.2 • D ✓
• C ✓ (Any 1) (1)
- 3.1.3 • B ✓
• E ✓ (Any 1) (1)

3.1.4 • **A** ✓ (1)

3.1.5 • **B** ✓ (1)

3.2 Temperature requirements

3.2.1 Heat production

- Dairy cows/cattle. ✓ (1)
- Reason**
- It has a value of 2500 kJ/h which is more than the others. ✓
 - Millions/many micro-organisms in the stomach that produce heat through fermentation. ✓ (Any 1) (1)

3.2.2 Reasons for intensive chicken production

- Heat production is the lowest ✓
- Optimal temperature is the highest ✓
- Lower critical temperature is the highest ✓
- Close range between critical and optimal temperature ✓
- Air movement to prevent respirational problems ✓
- Temperature control to prevent diseases ✓ (Any 2) (2)

3.2.3 Definition of optimal temperature

- Ideal/best/most comfortable/most suitable /most favourable environmental temperature ✓
- The animal does not need to use its own energy to control body temperature/allows for the most cost effective production output/and conducive for production ✓ (2)

3.2.4 Reason for keeping pigs in an enclosed environment in winter

- Have a higher lower critical temperature compared to cows ✓
- And will need a warmer environment to survive/to produce ✓
- As they have a lower ability to produce heat ✓ (Any 2) (2)

3.3 Adaptation of Mbuzi goat to harsh conditions

3.3.1 (a) Environmental conditions

- Pigmentation of the skin ✓
- protects it from radiation ✓
- or
- Legs adapted ✓
- to steeper slopes/to reach nutritious vegetation ✓
- or
- Horns ✓
- to protect from predators ✓
- or
- Good motherly instincts ✓
- supply it's young with regular nutrition (sucking of milk) ✓ (2)

(b) Pest and diseases

- Nutritious shrubs and bushes ✓
- makes it strong and more resistant to infections ✓
- or
- Not susceptible to worm/internal parasites ✓
- due to adaptability to local condition ✓
- or
- Indigenous breed which has been exposed to local conditions including pests and diseases ✓
- That is naturally selected to be resistant ✓ (2)

3.3.2 Management practices

- Dipping/pest control ✓
- Injection/Vaccination ✓
- Counting ✓
- Dosing/Deworming ✓
- Weaning/Kidding ✓
- Tagging/marking/identification ✓
- Castration ✓
- Weighting ✓
- Age determination ✓
- Artificial Insemination/AI ✓
- Hoof trimming ✓ (Any 3) (3)

3.4 Feed utilisation and conversion

3.4.1 Calculation of feed for Animal A and B

- Quantity of feed for **Animal A**: $48 \text{ kg} - 6 \text{ kg} = 42 \text{ kg}$ ✓
- Quantity of feed for **Animal B**: $56 \text{ kg} - 6 \text{ kg} = 50 \text{ kg}$ ✓ (2)

3.4.2 Animal that used the ration more effectively

- Animal B ✓

(1)

Reason

- Higher production in proportion to feed ingested ✓
- Animal A : 42kg of feed to produce 12 litres of milk/3.5kg per litre
Animal B : 50kg of feed to produce 43 litres of milk/ 1,2kg per litre ✓
- Better conversion rate ✓
- Better genetic material ✓

(Any 2) (2)

3.5 Broiler production unit**3.5.1 Equipment to be used (each equipment should be used once)**

- Insulation material on the roof/foldable walls ✓
- Electric heaters/foldable walls ✓
- Fans on the roof and walls/ foldable walls ✓
- Foot bath at all entrances ✓

(1)

(1)

(1)

(1)

3.5.2 Characteristics of an intensive animal production

- It is highly hygienic ✓
- Protection against extreme environmental conditions/shelter is provided ✓
- Regulation of optimal temperature for production ✓
- Fixed/regular supply of feed and water ✓
- Kept at high density/lots of animals in a small space ✓
- Capital intensive/expensive with high tech equipment/mechanized ✓

(Any 4) (4)

3.5.3 Possible health risks

- Disease ✓
- The outbreak of an infectious disease may be transmitted quickly amongst poultry ✓

(2)

[35]**QUE/STION 4: ANIMAL REPRODUCTION, PROTECTION AND CONTROL****4.1 Pregnancy testing****4.1.1 Labelled parts**

A - Caruncles/uterus wall/placenta ✓

B - Uterus/womb ✓

C – Cervix ✓

(3)

4.1.2 TWO advantages of early pregnancy testing

- For proper feeding ✓
- For proper management of diseases and parasites ✓
- For proper management of breeding cycles/records/calving date ✓

(Any 2) (2)

4.1.3 Role of the mucus plug

- Protects animal ✓
- Against external diseases and infections ✓

(2)

4.1.4 Conditions that could occur

(a) Maceration ✓


(1)

(b) Mummification ✓

(1)

- 4.1.5 **Condition and TWO possible causes for the termination of pregnancy** (1)
- Abortion/miscarriage ✓
- Reason**
- Diseases ✓
 - Injuries/maltreatment ✓
 - Infection ✓
 - Laxatives/wrong medication ✓
 - Toxins/poisons ✓
 - Fever reaction ✓
 - Malnutrition ✓
 - Stress ✓
 - Hormonal imbalances ✓ (Any 2) (2)
- 4.2 **Structures in reproduction**
- 4.2.1 **Names represented by letters**
- A - Ovary ✓
B - Graafian follicle ✓
E - Infundibulum ✓ (3)
- 4.2.2 **Hormones**
- (a) Follicle stimulating hormone/FSH ✓ (1)
(b) Oestrogen/LH (Luteinising hormone) ✓ (1)
(c) Oestrogen ✓ (1)
(d) Progesterone ✓ (1)
- 4.2.3 **Adaptability of infundibulum**
- Contains hair-like structures/cilia ✓
 - for movement of the ova ✓
- Or**
- Wider at the edge/bell shaped/ funnel shape ✓
 - adapted for holding/capturing the ova ✓ (2)
- 4.3 **Indigenous methods of controlling diseases**
- 4.3.1 Herbs/concoctions/mixtures ✓ (1)
- 4.3.2 **TWO most common methods**
- Gall smearing ✓
 - Self diagnosis ✓ (2)
- 4.3.3 **Ways in which bush burning control the ticks**
- Destroy eggs/larva/nymph ✓
 - Kills adult ticks ✓
 - Host in the life cycle for 2 and 3 host ticks is killed ✓ (2)

4.3.4 **Bar graph on indigenous control methods**



Marking graph with the following checklist:

Criteria	Yes: 1 Mark	No: 0 Mark
1. Bar graph	1 ✓	
2. X axis labelled	1 ✓	
3. Y axis labelled	1 ✓	
4. Points are plotted correctly	1 ✓	
5. Correct heading	1 ✓	
6. Correct subheadings for X- axis	1 ✓	

(6)

4.3.5 **THREE measures to restrict infectious diseases**

- Vaccination/inoculation ✓
- Injections ✓
- Bio-security/sanitation/proper handling of manure ✓
- Sufficient space/ good ventilation ✓
- Isolation/quarantine/separation ✓
- Controlling pests and parasites / dipping/ dosing ✓
- Good nutrition and supplements ✓
- Breeding of resistant animals ✓

(Any 3)

(3)

[35]

TOTAL SECTION B: 105
GRAND TOTAL: 150