



# basic education

Department:  
Basic Education  
**REPUBLIC OF SOUTH AFRICA**

## **NASIONALE SENIOR SERTIFIKAAT**

**GRAAD 12**

**WISKUNDE V2**

**MODEL 2014**

**MEMORANDUM**

**PUNTE: 150**

**Hierdie memorandum bestaan uit 13 bladsye.**



**VRAAG 2**

<p>2.1</p>		<p>✓ anker by 0 ✓ plot by boonste limiete ✓ gladde kurwe</p> <p>(3)</p>
<p>2.2</p>	<p><math>40 \leq t &lt; 60</math></p>	<p>✓ klas</p> <p>(1)</p>
<p>2.3</p>	<p>(96 ; 164) ∴ 172 – 164 = 8 leerdere</p>	<p>✓ 164 ✓ 8</p> <p>(2)</p>
<p>2.4</p>	<p>Frekwensie: 25; 44; 60; 28; 9; 6</p> $\text{gemiddelde} = \frac{25 \times 10 + 44 \times 30 + 60 \times 50 + 28 \times 70 + 9 \times 90 + 6 \times 110}{172}$ $= \frac{8000}{172}$ $= 46,51 \text{ uur}$	<p>✓ frekwensie ✓ middelpunte</p> <p>✓ <math>\frac{8000}{172}</math> ✓ antwoord</p> <p>(4) <b>[10]</b></p>

**VRAAG 3**


3.1	$K(7 ; 0)$	✓ antwoord (1)
3.2	$1 = \frac{x_M + 7}{2}$ en $1 = \frac{y_M + 3}{2}$ $\therefore M(-5 ; -1)$	✓ $x$ ✓ $y$ (2)
3.3	$m_{PM} = \frac{3-1}{7-1}$ $= \frac{1}{3}$	✓ substitusie ✓ antwoord (2)
3.4	$\tan \hat{P}\hat{S}\hat{K} = m_{PM} = \frac{1}{3}$ $\hat{P}\hat{S}\hat{K} = \tan^{-1}\left(\frac{1}{3}\right) = 18,43^\circ$ $\therefore \theta = 180^\circ - 90^\circ - 18,43^\circ = 71,57^\circ$	✓ $\tan \hat{P}\hat{S}\hat{K} = m_{PM}$ ✓ $\hat{P}\hat{S}\hat{K}$ ✓ $\theta$ (3)
3.5	$\cos 71,57^\circ = \frac{PK}{PS} = \frac{3}{PS}$ $PS = \frac{3}{\cos 71,57^\circ}$ $= 9,49$ eenhede <b>OF</b> $\sin 18,43^\circ = \frac{PK}{PS} = \frac{3}{PS}$ $PS = \frac{3}{\sin 18,43^\circ}$ $= 9,49$ eenhede	✓ korrekte verhouding ✓ PS onderwerp ✓ antwoord (3)  ✓ korrekte verhouding ✓ PS onderwerp ✓ antwoord (3)
3.6	$N(x ; -2x + 17)$ $m_{TN} = m_{PM}$ (TN    PM) $\frac{-2x + 17 - 5}{x - (-1)} = \frac{1}{3}$ $-6x + 36 = x + 1$ $-7x = -35$ $x = 5$ $\therefore y = -2(5) + 17 = 7$ $\therefore N(5 ; 7)$ <b>OF</b>	✓ N in terme van $x$ ✓ gelyke gradiënte ✓ substitusie  ✓ $x$ -waarde ✓ $y$ -waarde (5)

	$m_{TM} = \frac{1}{3} \quad (\text{TN} \parallel \text{PM})$ <p>vergelyking van TM:</p> $y - y_1 = \frac{1}{3}(x - x_1)$ $y - 5 = \frac{1}{3}(x - (-1))$ $y - 5 = \frac{1}{3}x + \frac{1}{3}$ $y = \frac{1}{3}x + 5\frac{1}{3}$ <p style="text-align: center;"><b>OF</b></p> $y = \frac{1}{3}x + c$ $5 = \frac{1}{3}(-1) + c$ $5\frac{1}{3} = c$ $y = \frac{1}{3}x + 5\frac{1}{3}$ $-2x + 17 = \frac{1}{3}x + 5\frac{1}{3}$ $-2\frac{1}{3}x = -11\frac{2}{3}$ $x = 5$ $\therefore y = -2(5) + 17 = 7$ $\therefore N(5; 7)$	<p>✓ <math>m_{TM}</math></p> <p>✓ vergelyking van TM</p> <p>✓ stel gelyk aan mekaar</p> <p>✓ <math>x</math>-waarde</p> <p>✓ <math>y</math>-waarde</p> <p style="text-align: right;">(5)</p>
<p>3.7.1</p>	<p><math>y = 5</math></p>	<p>✓ vergelyking</p> <p style="text-align: right;">(1)</p>
<p>3.7.2</p>	<p>gradiënt van AQ = <math>\tan 45^\circ</math> of <math>\tan 135^\circ</math>  <math>= 1</math> of <math>-1</math></p> $m_{AQ} = \frac{5-1}{a-1} = \pm 1$ $\therefore a - 1 = 4 \text{ of } -4$ $\therefore a = 5 \text{ of } -3$	<p>✓ <math>m_{AQ} = 1</math> of</p> <p>✓ <math>m_{AQ} = -1</math></p> <p>✓ substitusie in gradiëntformule</p> <p>✓ <math>x</math>-waarde</p> <p>✓ <math>y</math>-waarde</p> <p style="text-align: right;">(5)  <b>[22]</b></p>

**VRAAG 4**

4.1	$M(-1 ; -1)$	✓ antwoord (1)
4.2	$m_{NT} = \frac{2-1}{3-4} = -1$ $\therefore m_{AT} = 1 \quad (\text{radius} \perp \text{raaklyn})$ $y - 1 = 1(x - 4)$ $y = x - 3$	✓ $m_{NT}$ ✓ $m_{AT}$ ✓ rede ✓ substitusie van $m$ en $(4 ; 1)$ ✓ vergelyking (5)
4.3	$MR \perp AB$ (lyn vanaf midpt na midpt van koord) $MB^2 = MR^2 + RB^2$ (Stelling van Pythagoras) $9 = \left(\frac{\sqrt{10}}{2}\right)^2 + RB^2$ $RB^2 = \frac{13}{2}$ $RB = \sqrt{\frac{13}{2}}$ $AB = 2\left(\sqrt{\frac{13}{2}}\right) = \sqrt{26}$ eenhede	✓ $MR \perp AB$ ✓ $MB = 3$ ✓ substitusie in stelling van Pythagoras ✓ $AB$ in wortelvorm (4)
4.4	$MN^2 = (-1 - 3)^2 + (-1 - 2)^2$ $= 16 + 9$ $= 25$ $MN = 5 \text{ eenhede}$	✓ substitusie in afstandformule ✓ antwoord (2)
4.5	$r = 5 - 3 = 2$ eenhede $\therefore (x - 3)^2 + (y - 2)^2 = 4$ $\therefore x^2 + y^2 - 6x - 4y + 9 = 0$	✓ $r$ ✓ substitusie in sirkelvergelyking ✓ vergelyking (3) <b>[15]</b>

**VRAAG 5**

<p>5.1.1</p>	$-\sin \alpha$ $= -\left(-\frac{4}{5}\right) = \frac{4}{5}$	<p>✓ reduksie</p> <p>✓ antwoord</p> <p>(2)</p>
<p>5.1.2</p>	$(-4)^2 + b^2 = 5^2$ $b^2 = 25 - 16 = 9$ $b = -3$ $\cos \alpha = \frac{-3}{5}$ 	<p>✓ <math>b = -3</math></p> <p>✓ antwoord</p> <p>(2)</p>
<p>5.1.3</p>	$\sin(\alpha - 45^\circ)$ $= \sin \alpha \cos 45^\circ - \cos \alpha \sin 45^\circ$ $= -\frac{4}{5} \cdot \frac{1}{\sqrt{2}} - \left(-\frac{3}{5}\right) \cdot \frac{1}{\sqrt{2}}$ $= -\frac{1}{5\sqrt{2}}$ <p style="text-align: center;"><b>OF</b></p> $\sin(\alpha - 45^\circ)$ $= \sin \alpha \cos 45^\circ - \cos \alpha \sin 45^\circ$ $= -\frac{4}{5} \cdot \frac{\sqrt{2}}{2} - \left(-\frac{3}{5}\right) \cdot \frac{\sqrt{2}}{2}$ $= -\frac{\sqrt{2}}{10}$	<p>✓ uitbreiding</p> <p>✓ <math>\frac{1}{\sqrt{2}}</math></p> <p>✓ antwoord in eenvoudigste vorm</p> <p>(3)</p> <p>✓ uitbreiding</p> <p>✓ <math>\frac{\sqrt{2}}{2}</math></p> <p>✓ antwoord in eenvoudigste vorm</p> <p>(3)</p>
<p>5.2.1</p>	$LHS = \frac{8 \sin x \cdot \cos x}{\sin^2 x - \cos^2 x}$ $= \frac{4(2 \sin x \cdot \cos x)}{\sin^2 x - \cos^2 x}$ $= \frac{4 \sin 2x}{-(\cos^2 x - \sin^2 x)}$ $= \frac{4 \sin 2x}{-\cos 2x}$ $= -4 \tan 2x$	<p>✓ <math>\sin x</math></p> <p>✓ <math>\cos x</math></p> <p>✓ <math>\cos^2 x</math></p> <p>✓ <math>4 \sin 2x</math></p> <p>✓ faktoriseer</p> <p>✓ <math>-\cos 2x</math></p> <p>(6)</p>
<p>5.2.2</p>	<p>Ongedefinieer as <math>\cos 2x = 0</math> of <math>\tan 2x = \infty</math>:</p> <p><math>x = 45^\circ</math> en</p> <p><math>x = 135^\circ</math></p>	<p>✓ <math>45^\circ</math></p> <p>✓ <math>135^\circ</math></p> <p>(2)</p>


5.3	$1 - 2\sin^2 \theta + 4\sin^2 \theta - 5\sin \theta - 4 = 0$ $2\sin^2 \theta - 5\sin \theta - 3 = 0$ $(2\sin \theta + 1)(\sin \theta - 3) = 0$ <p><math>\therefore \sin \theta = -\frac{1}{2}</math> of <math>\sin \theta = 3</math> (geen oplossing)</p> <p><math>\therefore \theta = 210^\circ + 360^\circ k</math> of <math>\theta = 330^\circ + 360^\circ k ; k \in \mathbb{Z}</math></p> <p><b>OF</b></p> <p><math>\therefore \theta = 210^\circ + 360^\circ k</math> of <math>\theta = 30^\circ + 360^\circ k ; k \in \mathbb{Z}</math></p>	<ul style="list-style-type: none"> <li>✓ <math>1 - 2\sin^2 \theta</math></li> <li>✓ standaardvorm</li> <li>✓ faktore</li> <li>✓ geen oplossing</li> <li>✓ <math>210^\circ</math></li> <li>✓ <math>330^\circ</math></li> <li>✓ <math>+ 360^\circ k ; k \in \mathbb{Z}</math></li> </ul> <p style="text-align: right;">(7) <b>[22]</b></p>
-----	---	--

**VRAAG 6**

6.1	$b = \frac{1}{2}$	✓ waarde van $b$ (1)
6.2	A( $30^\circ ; 1$ )	✓ $30^\circ$ ✓ 1 (2)
6.3	$x = 160^\circ$	✓ $x = 160^\circ$ (1)
6.4	$h(x) = 2\cos(x - 30^\circ) + 1$ $y \in [-1 ; 3]$ <b>OF</b> $-1 \leq y \leq 3$	<ul style="list-style-type: none"> <li>✓ kritiese waardes</li> <li>✓ notasie</li> </ul> <p style="text-align: right;">(2) <b>[6]</b></p>



**VRAAG 7**

<p>7.1</p>	<p>Trek <math>CD \perp AB</math>                  In <math>\triangle ACD</math>:  <math>\sin A = \frac{CD}{b} \therefore CD = b \cdot \sin A</math>                   In <math>\triangle CBD</math>:  <math>\sin B = \frac{CD}{a} \therefore CD = a \cdot \sin B</math>   <math>\therefore b \cdot \sin A = a \cdot \sin B</math>  <math>\therefore \frac{\sin A}{a} = \frac{\sin B}{b}</math></p> 	<p>✓ konstruksie                  ✓ sin A                  ✓ maak CD die onderwerp                   ✓ sin B                   ✓ <math>b \cdot \sin A = a \cdot \sin B</math>                  (5)</p>
<p>7.2.1</p>	<p><math>\widehat{SPQ} = 180^\circ - 2x</math> (teenoorst <math>\angle e</math> van koordevierh )  <math>\widehat{PSQ} + \widehat{PQS} = 2x</math> (som van <math>\angle e</math> in <math>\triangle</math>)  <math>\widehat{PSQ} = \widehat{PQS} = x</math> (<math>\angle e</math> teenoor gelyke sye)</p>	<p>✓ <math>\widehat{SPQ} = 180^\circ - 2x</math> (S/R)                  ✓ rede                  (2)</p>
<p>7.2.2</p>	$\frac{\sin \widehat{SPQ}}{\sin(180^\circ - 2x)} = \frac{\sin \widehat{PSQ}}{\sin x}$ $\frac{SQ}{\sin(180^\circ - 2x)} = \frac{PQ}{\sin x}$ $SQ = \frac{k \sin 2x}{\sin x}$ $SQ = \frac{k(2 \sin x \cdot \cos x)}{\sin x} = 2k \cos x$ <p style="text-align: center;"><b>OF</b></p> $SQ^2 = PQ^2 + PS^2 - 2PQ \cdot PS \cdot \cos \widehat{SPQ}$ $= k^2 + k^2 - 2 \cdot k \cdot k \cdot \cos(180^\circ - 2x)$ $= 2k^2 + 2k^2 \cos 2x$ $= 2k^2 + 2k^2(2\cos^2 x - 1)$ $= 4k^2 \cos^2 x$ $SQ = 2k \cos x$	<p>✓ substitusie in korrekte formule                  ✓ sin 2x                   ✓ SQ onderwerp                  ✓ <math>2 \sin x \cdot \cos x</math>                  (4)                   ✓ substitusie in korrekte formule                  ✓ <math>-\cos 2x</math>                  ✓ <math>2\cos^2 x - 1</math>                  ✓ vereenvoudig                  (4)</p>
<p>7.2.3</p>	$\tan y = \frac{3}{k}$ $k = \frac{3}{\tan y}$ $SQ = 2 \cos x \left( \frac{3}{\tan y} \right)$ $\therefore = \frac{6 \cos x}{\tan y}$	<p>✓ tan-verhouding                   ✓ <math>k</math> onderwerp en substitusie                  (2)  <b>[13]</b></p>

**VRAAG 8**

8.1	die hoek onderspan in die teenoorstaande sirkelsegment	✓korrekte stelling (1)
8.2.1	$\hat{B}_1 = \hat{E}_1 = 68^\circ$ (rkl-koordst)	✓ $\hat{E}_1 = 68^\circ$ ✓ rede (2)
8.2.2	$\hat{E}_1 = \hat{B}_3 = 68^\circ$ (verwiss $\angle$ e; AE    BC)	✓ $\hat{B}_3 = 68^\circ$ (S/R) (1)
8.2.3	$\hat{D}_1 = \hat{B}_3 = 68^\circ$ (buite $\angle$ v koordevh)	✓ $\hat{D}_1 = 68^\circ$ ✓ rede (2)
8.2.4	$\hat{E}_2 = 20^\circ + 68^\circ$ $= 88^\circ$ (buite $\angle$ v $\Delta$ )	✓ $\hat{E}_2 = 88^\circ$ (S/R) (1)
8.2.5	$\hat{C} = 180^\circ - 88^\circ$ $= 92^\circ$ (tos $\angle$ e v koordevh)	✓ $\hat{C} = 92^\circ$ ✓ rede (2) <b>[9]</b>

**VRAAG 9**

<p>9.1</p>	<p><math>\hat{D}_4 = \hat{A} = x</math> (rkl-koordstelling)   <math>\hat{A} = \hat{D}_2 = x</math> (<math>\angle</math>e tos gelyke sye)</p>	<p>✓ <math>\hat{A} = x</math>                  ✓ rede                  ✓ <math>\hat{A} = \hat{D}_2 = x</math> (S/R)                  (3)</p>
<p>9.2</p>	<p><math>\hat{M}_1 = 2x</math> (buite <math>\angle</math>v<math>\Delta</math>) OF (<math>\angle</math> by midpt = 2<math>\angle</math> by omtr)  <math>\hat{M}\hat{D}E = 90^\circ</math> (radius <math>\perp</math> rkl)  <math>\hat{M}_2 = 90^\circ - 2x</math>  <math>\therefore \hat{E} = 180^\circ - (90^\circ + 90^\circ - 2x)</math> (som v <math>\angle</math>e in <math>\Delta</math>MDE)  <math>= 2x</math>  <math>\therefore</math> CM is 'n rkl (omgek rkl-koordst)</p>	<p>✓ <math>\hat{M}_1 = 2x</math> (S/R)                  ✓ <math>\hat{M}\hat{D}E = 90^\circ</math> (S/R)                  ✓ <math>\hat{E} = 2x</math>                  ✓ rede                  (4)</p>
<p>9.3</p>	<p><math>\hat{M}_3 = 90^\circ</math> (EM <math>\perp</math> AC)  <math>\hat{A}\hat{D}B = 90^\circ</math> (<math>\angle</math> in halfsirkel)  <math>\therefore</math> FMBD is koordevh (buite<math>\angle</math> v vh = tos binne <math>\angle</math>)  <b>OF</b>  <math>\hat{E}\hat{M}C = 90^\circ</math> (EM <math>\perp</math> AC)  <math>\hat{A}\hat{D}B = 90^\circ</math> (<math>\angle</math> in halfsirkel)  <math>\therefore</math> FMBD is koordevh (tos <math>\angle</math>e v vh suppl)</p>	<p>✓ <math>\hat{M}_3 = 90^\circ</math>                  ✓ <math>\hat{A}\hat{D}B = 90^\circ</math> (S/R)                  ✓ rede (3)                  ✓ <math>\hat{E}\hat{M}C = 90^\circ</math>                  ✓ <math>\hat{A}\hat{D}B = 90^\circ</math> (S/R)                  ✓ rede (3)</p>
<p>9.4</p>	<p><math>DC^2 = MC^2 - MD^2</math> (Pythagoras)  <math>= (3BC)^2 - (2BC)^2</math> (MB = MD = radii)  <math>= 9BC^2 - 4BC^2</math>  <math>= 5BC^2</math></p>	<p>✓ Pythagoras                  ✓ substitusie                  ✓ <math>9BC^2 - 4BC^2</math> (3)</p>
<p>9.5</p>	<p>In <math>\Delta</math>DBC en <math>\Delta</math>DFM:  <math>\hat{D}_4 = \hat{D}_2 = x</math> (bewys in 9.1)  <math>\hat{B}_1 = \hat{F}_2</math> (buite <math>\angle</math> v koordevh)  <math>\hat{C} = \hat{M}_2</math>  <math>\therefore \Delta</math>DBC <math>\parallel\parallel\parallel</math> <math>\Delta</math>DFM (<math>\angle</math>; <math>\angle</math>; <math>\angle</math>)</p>	<p>✓ <math>\hat{D}_4 = \hat{D}_2</math>                  ✓ <math>\hat{B}_1 = \hat{F}_2</math>                  ✓ rede                  ✓ <math>\hat{C} = \hat{M}_2</math> of (<math>\angle</math>; <math>\angle</math>; <math>\angle</math>) (4)</p>
<p>9.6</p>	<p><math>\frac{DM}{FM} = \frac{DC}{BC}</math> (<math>\Delta</math>DBC <math>\parallel\parallel\parallel</math> <math>\Delta</math>DFM)  <math>= \frac{\sqrt{5}BC}{BC}</math>  <math>= \sqrt{5}</math></p>	<p>✓ S                  ✓ antwoord (2)  <b>[19]</b></p>

**VRAAG 10**

<p>10.1</p>		<p>✓ konstruksie</p> <p>✓ <math>\frac{\text{opp } \triangle ADE}{\text{opp } \triangle DEB} = \frac{AD}{DB}</math></p> <p>✓ rede</p> <p>✓ <math>\frac{\text{opp } \triangle ADE}{\text{opp } \triangle DEC} = \frac{AE}{EC}</math></p> <p>✓ Area <math>\triangle DEB =</math> Area <math>\triangle DEC</math> (S/R)</p> <p>✓</p> <p><math>\frac{\text{opp } \triangle ADE}{\text{opp } \triangle DEB} = \frac{\text{opp } \triangle ADE}{\text{opp } \triangle DEC}</math></p>
	<p>Konstruksie: Verbind DC en BE en trek hoogtes <math>k</math> en <math>h</math></p> <p><math>\frac{\text{opp } \triangle ADE}{\text{opp } \triangle DEB} = \frac{\frac{1}{2} \cdot AD \cdot k}{\frac{1}{2} \cdot DB \cdot k} = \frac{AD}{DB}</math> (gelyke hoogtes)</p> <p><math>\frac{\text{opp } \triangle ADE}{\text{opp } \triangle DEC} = \frac{\frac{1}{2} \cdot AE \cdot h}{\frac{1}{2} \cdot EC \cdot h} = \frac{AE}{EC}</math> (gelyke hoogtes)</p> <p>Maar Opp <math>\triangle DEB =</math> Opp <math>\triangle DEC</math> (dies basis, dies hoogte)</p> <p><math>\therefore \frac{\text{opp } \triangle ADE}{\text{opp } \triangle DEB} = \frac{\text{opp } \triangle ADE}{\text{opp } \triangle DEC}</math></p> <p><math>\therefore \frac{AD}{DB} = \frac{AE}{EC}</math></p>	<p>(6)</p>

<p>10.2.1</p>	$\frac{AB}{BE} = \frac{AC}{CD}$ <p>(Ewered st; BC    ED)</p> $\frac{1}{3} = \frac{3}{CD}$ $\therefore CD = 9 \text{ eenhede}$	<p>✓ <math>\frac{AB}{BE} = \frac{AC}{CD}</math> (S/R) ✓ substitusie ✓ antwoord (3)</p>
<p>10.2.2</p>	$\frac{DG}{GA} = \frac{FD}{FE}$ <p>(Ewered st; FG    EA)</p> $\frac{9-x}{3+x} = \frac{3}{6}$ $54 - 6x = 9 + 3x$ $-9x = -45$ $x = 5$	<p>✓ <math>\frac{DG}{GA} = \frac{FD}{FE}</math> (S/R) ✓ substitusie ✓ vereenvoudig ✓ antwoord (4)</p>
<p>10.2.3</p>	<p>In <math>\triangle ABC</math> en <math>\triangle AED</math>:</p> <p><math>\hat{A}</math> is gemeen  <math>\hat{A}\hat{B}C = \hat{E}</math> (ooreenk <math>\angle</math>s; BC    ED)  <math>\hat{A}\hat{C}B = \hat{D}</math> (ooreenk <math>\angle</math>s; BC    ED)  <math>\triangle ABC \sim \triangle AED</math> (<math>\angle, \angle, \angle</math>)  <math>\therefore \frac{BC}{ED} = \frac{AC}{AD}</math>  <math>\frac{BC}{9} = \frac{3}{12}</math>  <math>BC = 2\frac{1}{4}</math> eenhede</p>	<p>✓ <math>\hat{A}</math> is gemeen          ✓ <math>\hat{A}\hat{B}C = \hat{E}</math> (S/R)          ✓ <math>\hat{A}\hat{C}B = \hat{D}</math> (S/R)          of (<math>\angle; \angle; \angle</math>)          ✓ <math>\frac{BC}{ED} = \frac{AC}{AD}</math>          ✓ antwoord (5)</p>
<p>10.2.4</p>	$\frac{\text{opp } \triangle ABC}{\text{opp } \triangle GFD} = \frac{\frac{1}{2} AC \cdot BC \cdot \sin \hat{A}\hat{C}B}{\frac{1}{2} GD \cdot FD \cdot \sin \hat{D}}$ $= \frac{\frac{1}{2} (3)(2\frac{1}{4}) \sin \hat{D}}{\frac{1}{2} (4)(3) \sin \hat{D}}$ <p>(ooreenk <math>\angle</math>s; BC    ED)</p> $= \frac{9}{16}$	<p>✓ gebruik v opp reël          ✓ korrekte sye en <math>\angle</math>e          ✓ substitusie v waardes          ✓ <math>\sin \hat{A}\hat{C}B = \sin \hat{D}</math> (S/R)          ✓ antwoord (5)  <b>[23]</b></p>

**TOTAAL: 150**